

SKRIPSI

IKE TERYASTUTIK

**PENGARUH KEPOLARAN PELARUT
PADA SINTESIS OKTIL 3-HIDROKSISINAMAT
DARI BAHAN AWAL ASAM 3-HIDROKSISINAMAT**



**FAKULTAS FARMASI UNIVERSITAS AIRLANGGA
BAGIAN KIMIA FARMASI
SURABAYA
2003**

Lembar Pengesahan

**PENGARUH KEPOLARAN PELARUT
PADA SINTESIS OKTIL 3-HIDROKSISINAMAT
DARI BAHAN AWAL ASAM 3-HIDROKSISINAMAT**

SKRIPSI

**Dibuat Untuk Memenuhi Syarat Mencapai Gelar Sarjana Farmasi Pada
Fakultas Farmasi Universitas Airlangga
2003**

Oleh :

**IKE TERYASTUTIK
NIM : 059912118**

Disetujui Oleh :


Dra. Juni Ekowati, Msi., Apt
Pembimbing Utama


Dr. Tutuk Budiati, MS., Apt
Pembimbing Serta

ABSTRACT

The aim of this research is to compare the effect of polarity of solvent in the synthesis of octyl-3-hydroxycinnamate. The solvent used in this research were tetrahydrofuran (THF) and benzena. Tetrahydrofuran (THF) has dielectric constant 7,58 and dipole moment 1,74, while benzena has dielectric constant 2,28 and dipole moment 0. The resulted compound in tetrahydrofuran (THF) was liquid in room temperature and the colour was brownish yellow. It gave 49,43 % yields. The identification of the resulted compound was done by TLC test, refractive index, UV-Vis spectrophotometry, FT/IR spectrophotometry and gas chromatography-mass spectra (GC-MS).

There were no resulted compound in benzena. 3-hydroxycinnamic acid as starting material was not solvated in benzena so the ionization can't occur and octyl 3-hydroxycinnamate can't be resulted.

Key word : esterification reaction, benzena, tetrahydrofuran, octyl 3-hydroxycinnamate